arXiv:1602.01617v1 [math.CO] 4 Feb 2016

# The intricate labyrinth of Collatz Sequences

**Maya Mohsin Ahmed**

**Abstract**

In a previous article, we reduced the unsolved problem of the convergence of Collatz sequences, to convergence of Collatz sequences of odd numbers, that are divisible by 3. In this article, we further reduce this set to odd numbers that are congruent to 21 mod 24. We also show that either the Collatz sequence of a given odd number or an equivalent Collatz sequence reverses to a multiple of 3. Moreover, we construct a network composed of Collatz sequences of all odd numbers.

## 1 Introduction.

Given a positive integer $A$, construct the sequence $c_i$ as follows:

$$\begin{aligned} c_i &= A && \text{if } i = 0; \\ &= 3c_{i-1} + 1 && \text{if } c_{i-1} \text{ is odd;} \\ &= c_{i-1}/2 && \text{if } c_{i-1} \text{ is even.} \end{aligned}$$

The sequence $c_i$ is called a *Collatz sequence* with *starting number* $A$. The *Collatz conjecture* says that this sequence will eventually reach the number 1, regardless of which positive integer is chosen initially. The sequence gets in to an infinite cycle of 4, 2, 1 after reaching 1.

**Example 1.1.** The Collatz sequence of 27 is:

$27, 82, 41, 124, 62, 31, 94, 47, 142, 71, 214, 107, 322, 161, 484, 242, 121, 364, 182, 91, 274, 137,$
$412, 206, 103, 310, 155, 466, 233, 700, 350, 175, 526, 263, 790, 395, 1186, 593, 1780, 890, 445,$
$1336, 668, 334, 167, 502, 251, 754, 377, 1132, 566, 283, 850, 425, 1276, 638, 319, 958, 479, 1438,$
$719, 2158, 1079, 3238, 1619, 4858, 2429, 7288, 3644, 1822, 911, 2734, 1367, 4102, 2051, 6154,$
$3077, 9232, 4616, 2308, 1154, 577, 1732, 866, 433, 1300, 650, 325, 976, 488, 244, 122, 61, 184, 92,$
$46, 23, 70, 35, 106, 53, 160, 80, 40, 20, 10, 5, 16, 8, 4, 2, 1, 4, 2, 1, 4, 2, 1, \ldots$

For the rest of this article, we will ignore the infinite cycle of $4, 2, 1$, and say that a Collatz sequence *converges to* 1, if it reaches 1. The Collatz conjecture is an unsolved conjecture about the convergence of a sequence named after Lothar Collatz. A comprehensive study of the Collatz conjecture can be found in [2], [3], and [4].

In this article, like in [1], we focus on the subsequence of odd numbers of a Collatz sequence. This is because every even number in a Collatz sequence has to reach an odd

number after a finite number of steps. Observe that the Collatz conjecture implies that the subsequence of odd numbers of a Collatz sequence converges to 1.

**Example 1.2.** The subsequence of odd numbers of the Collatz sequence of 27 in Example 1.1 is:

$$27, 41, 31, 47, 71, 107, 161, 121, 91, 137, 103, 155, 233, 175, 263, 395, 593, 445, 167, 251, 377, 283, 425, 319, 479, 719, 1079, 1619, 2429, 911, 1367, 2051, 3077, 577, 433, 325, 61, 23, 35, 53, 5, 1, 1, \ldots$$

If two Collatz sequences become equal after a finite number of terms, we say they are *merging sequences*. We say two Collatz sequences are *equivalent* if the second odd number occurring in the sequences are same.

Let $A$ be an odd number. If $A > 1$, write $A = 2^k N + 1$ such that $N$ is an odd number and $k \geq 1$. In Section 2, we prove if $k = 2$ then Collatz sequences of $A$ and $N$ are equivalent; if $k > 2$ then Collatz sequences of $A$ and $2A + 1$ are equivalent; and finally, if $k = 1$, then the Collatz sequence of $A$ either merges with the Collatz sequence of $N$ or $2A + 1$.

Let $A$ be a positive integer. In [1], we defined a sequence in the reverse direction defined as follows. The *reverse Collatz sequence* $r_i$ of $A$ is given by

$$r_i = \begin{cases} A & \text{if } i = 0; \\ \frac{r_{i-1}-1}{3} & \text{if } r_{i-1} \equiv 1 \bmod 3 \text{ and } r_{i-1} \text{ is even}; \\ 2r_{i-1} & \text{if } r_{i-1} \not\equiv 1 \bmod 3 \text{ and } r_{i-1} \text{ is even}; \\ 2r_{i-1} & \text{if } r_{i-1} \text{ is odd}. \end{cases}$$

We say that a reverse Collatz sequence *converges* if the subsequence of odd numbers of the sequence converges to a multiple of 3. Let $A$ be an odd number. We showed in [1] (also see Section 2), that the Collatz sequences of $A$ and $4A + 1$ are equivalent. In [1], we conjectured that the reverse Collatz sequence of every number converges. Though, we cannot prove that conjecture yet, in Section 3, we show that the reverse Collatz sequence of either $A$ or $4A + 1$ converges. We also discuss how the convergence of the reverse Collatz sequences of $A$, $2A + 1$, and $4A + 1$ are related in Section 3.

Given an odd number $A$, in Section 4, we show that the Collatz sequence of $A$ is equivalent to an odd number of the form $24t + 21$, where $t \geq 0$. This result reduces the unsolved problem of convergence of Collatz sequences to the set of odd numbers that are divisible by 3 and are congruent to 21 modulo 24.

Finally, in Section 5, we construct a network of diagonal arrays which contain the Collatz sequences of all odd numbers.

## 2 The coupling of the Collatz sequences of $A$ and $2A+1$.

In this section, we discuss the merging of the Collatz sequence of an odd number $N$ with the Collatz sequences of $2N+1$ and $4N+3$.

If $A = 4^d \times P + 4^{d-1} + 4^{d-2} + \cdots + 4 + 1$, where $P$ is an integer and $d \geq 1$, then we say that $A$ is a *jump from $P$ of height $d$*. If $P \not\equiv 1 \bmod 4$, then $d$ is the *maximum height* of $A$.

**Example 2.1.** $53 = 4 \times 13 + 1 = 4^2 \times 3 + 4 + 1$ is a jump from 13 of height 1 and a jump from 3 of height 2.

**Proposition 2.1.** *Let $A$ be an odd number and let $a_i$ denote the sequence of odd numbers in the Collatz sequence of $A$ with $a_0 = A$.*

1. *Consider the sequence $c_i = 4c_{i-1} + 1$ with $c_0 = A$. Then for any $i$, the Collatz sequence of $A$ and $c_i$ are equivalent.*
2. *For some $r \geq 0$, $a_r \equiv 1 \bmod 4$, and*
$$a_i = \frac{3a_{i-1}+1}{2}, i = 1, \ldots, r, \text{ and } a_{r+1} = \frac{3a_r+1}{2^k}, \text{ for some } k \geq 2.$$
3. *If $A \equiv 1 \bmod 4$, and if $A > 1$, then $A = 2^k \times N + 1$ for some odd number $N$ and $k \geq 2$. Let $n_i$ denote the odd numbers in the Collatz sequence of $N$ with $n_0 = N$.*
   (a) *If $k = 2$, then $a_1 = n_1$. Moreover, if $A = 4^d P + 4^{d-1} + \cdots + 4^2 + 4 + 1$, such that $P \not\equiv 1 \bmod 4$, then*
   $$a_1 = \frac{3A+1}{4^d \times 2^i} = \frac{3P+1}{2^i},$$
   *where $i = 1$ if $P$ is odd, and $i = 0$ otherwise.*
   (b) *If $k > 2$, write $k = 2j + r$, $r = 0$ or $r = 1$. If $r = 0$ then $a_i = 2^{k-2i} \times 3^i N + 1$, for $i = 1, \ldots, j-1$, and the Collatz sequence of $A$ merges with the Collatz sequence of $3^{j-1}N$. On the other hand, if $r = 1$, then $a_i = 2^{k-2i} \times 3^i N + 1$, for $i = 1, \ldots, j$.*

*Proof.*

1. This result is proved as Corollary 2 in [1].
2. We get this result by applying Theorem 2 in [1].
3. (a) When $k = 2$, $A = 4N + 1$, therefore $a_1 = n_1$, by part (1). Since $A$ is a jump from $P$ of height $d$, Lemma 2 in [1] implies $3A+1 = 4^d(3P+1)$. Consequently, $a_1 = (3A+1)/(4^d \times 2^i) = (3P+1)/2^i$. Since $P \not\equiv 1 \bmod 4$, if $P$ is odd, $P \equiv 3 \bmod 4$, which implies $3P + 1 \equiv 2 \bmod 4$. Consequently, $i = 1$. On the other hand, if $P$ is even, $3P + 1$ is odd, hence $i = 0$.

(b) When $k > 2$

$$\frac{3A+1}{4} = \frac{3 \times 2^k N + 4}{4} = 3 \times 2^{k-2} N + 1$$

is an odd number. Hence $a_1 = 2^{k-2} \times 3 \times N + 1$. We repeat this argument to get that if $r = 0$ then $a_i = 2^{k-2i} \times 3^i N + 1$, for $i = 1, \ldots, j-1$. Thus $a_{j-1} = 4 \times 3^{j-1} N + 1$. Consequently, by applying Part 1, we derive that the Collatz sequence of $A$ merges with the Collatz sequence of $3^{j-1}N$. Similarly, we get that if $r = 1$, then $a_i = 2^{k-2i} \times 3^i N + 1$, for $i = 1, \ldots, j$.

□

**Theorem 2.1.** *Let $N$ be an odd number. Let $n_0 = N$, $m_0 = 2n_0 + 1 = 2N + 1$, and $l_0 = 2m_0 + 1 = 4N+3$ . Let $n_i, m_i$, and $l_i$ denote the subsequence of odd numbers in the Collatz sequence of $n_0, m_0$, and $l_0$, respectively. Then, for some integer $r$, $n_{r+1} = (3n_r + 1)/2^k$ such that $k > 1$. Let $r$ be the smallest such integer. Then, $m_{r+2} = (3m_{r+1} + 1)/2^j$ for some $j > 1$, and*

$$\begin{aligned} &m_i = 2n_i + 1, \text{ for } i \leq r, \\ &m_{r+1} = 2^k n_{r+1} + 1 \\ &l_i = 2m_i + 1 \text{ for } i \leq r+1, \\ &l_{r+2} = 2^j m_{r+2} + 1 \end{aligned}$$

*If $k = 2$, then $m_i = n_i$ for $i > r + 1$. Otherwise, if $k > 2$ then $l_{r+2} = 4m_{r+2} + 1$ and $l_i = m_i$ for $i > r + 2$.*

*Proof.* By part (2) of Proposition 2.1, there exists an integer $r$, such that $n_{r+1} = (3n_r + 1)/2^k$ where $k > 1$. Since $r$ is the smallest such number, we have $n_i = \frac{3n_{i-1}+1}{2}$ for $0 \leq i \leq r$. Therefore, substituting $n_1 = (3n_0 + 1)/2$ and $m_0 = 2n_0 + 1$, we get

$$\frac{3m_0+1}{2} = \frac{3(2n_0+1)+1}{2} = 3n_0 + 2 = 2\left(\frac{3n_0+1}{2}\right) + 1 = 2n_1 + 1.$$

Now $(3m_0 + 1)/2$ is odd because $2n_1 + 1$ is an odd number. Therefore, $m_1 = (3m_0 + 1)/2 = 2n_1 + 1$. Repeating the above argument we prove that $m_i = 2n_i + 1$ for $i \leq r$.

Since

$$\begin{aligned} 3m_r + 1 &= 3(2n_r + 1) + 1 \text{ (substituting } m_r = 2n_r + 1) \\ &= 2(3n_r + 1) + 2 \text{ (rearranging terms)} \\ &= 2(2^k n_{r+1}) + 2 \text{ (substituting } 3n_r + 1 = 2^k n_{r+1}) \end{aligned}$$

we get $(3m_r + 1)/2 = 2^k n_{r+1} + 1$ is an odd number. Therefore $m_{r+1} = (3m_r + 1)/2 = 2^k n_{r+1} + 1$. Consequently, $(3m_{r+1} + 1)$ is divisible by 4, hence $m_{r+2} = (3m_{r+1} + 1)/2^j$ such that $j > 1$.

If $k = 2$, $m_{r+1} = 4n_{r+1} + 1$, hence $m_i = n_i$ for $i > r + 1$ by part (1) of Proposition 2.1.

Again, since $l_0 = 2m_0 + 1$ and $m_i = (3m_{i-1} + 1)/2$ for $i \leq r + 1$, we repeat the above arguments to conclude that $l_i = 2m_i + 1$ for $0 \leq i \leq r + 1$.

Since

$$\frac{3l_{r+1} + 1}{2} = \frac{3(2m_{r+1} + 1) + 1}{2} = 3m_{r+1} + 2,$$

and $3m_{r+1} + 2$ is an odd number, we get

$$l_{r+2} = \frac{3l_{r+1} + 1}{2} = 3m_{r+1} + 2. \tag{1}$$

Now

$$\frac{3m_{r+1} + 1}{4} = \frac{3(2^k n_{r+1} + 1) + 1}{4} = 3 \times 2^{k-2} n_{r+1} + 1.$$

If $k > 2$, $3 \times 2^{k-2} n_{r+1} + 1$ is odd. Therefore $m_{r+2} = \frac{3m_{r+1}+1}{4}$. Consequently,

$$4m_{r+2} + 1 = 3m_{r+1} + 2. \tag{2}$$

From equations 1 and 2, we get $l_{r+2} = 4m_{r+2} + 1$. Again, by part (1) of Proposition 2.1, we get $l_i = m_i$, for $i > r + 2$. □

**Example 2.2.** We illustrate Theorem 2.1 with an example. Let $A = 3$, $n_0 = A = 3$ $m_0 = 2n_0 + 1 = 7$, and $l_0 = 2m_0 + 1 = 15$. Let $n_i, m_i, l_i$ denote the odd numbers in the Collatz sequence of $n_0$, $m_0$, and $l_0$, respectively. Thus

$$\begin{aligned} &n_i : 3, 5, 1 \\ &m_i : 7, 11, 17, 13, 5, 1 \\ &l_i : 15, 23, 35, 53, 5, 1 \end{aligned}$$

Since $n_2 = \frac{3n_1+1}{2^4} = 1$, by Theorem 2.1,

$$\begin{aligned} &m_i = 2n_i + 1 \text{ for } 0 \leq i \leq 1, \\ &m_2 = 2^4 n_2 + 1 = 17 \\ &l_i = 2m_i + 1 \text{ for } 0 \leq i \leq 2, \\ &l_3 = 4m_3 + 1 = 53, \text{ and} \\ &l_i = m_i \text{ for } i \geq 4. \end{aligned}$$

Thus, in this case, the Collatz sequence of 15 merges with the Collatz sequence of 7.

We conclude this section by observing the following. If $A > 1$ is an odd number, then write $A = 2^k N + 1$, such that, $N$ is an odd number and $k \geq 1$. If $k = 2$ then, by Part 1 of Proposition 2.1, the Collatz sequences of $A$ and $N$ are equivalent. If $k > 2$, then the odd number $p$ that occurs after $A$, in the Collatz sequence of $A$, is $p = (3A + 1)/4$. This implies, by Theorem 2.1, that the number occurring after $2A + 1$, in the Collatz sequence of $2A + 1$, is $4p + 1$. Consequently, the Collatz sequences of $A$ and $2A + 1$ are equivalent. Finally, if $k = 1$, then, by Theorem 2.1, the Collatz sequence of $A$ either merges with the Collatz sequence of $N$ or $2A + 1$.

# 3 The collaborative convergence of the reverse Collatz sequences of $A$, $4A+1$, and $2A+1$.

Let $A$ be an odd number. In this section, we show that, if $A \equiv 0 \bmod 3$, then either the reverse Collatz sequence of $2A+1$ or $4A+1$ converge. If $A \equiv 1 \bmod 3$, then $2A+1 \equiv 0 \bmod 3$, and either the reverse Collatz sequence of $A$ or $4A+1$ converge. Finally, If $A \equiv 2 \bmod 3$, then $4A+1 \equiv 0 \bmod 3$, and either the reverse Collatz sequence of $A$ or $2A+1$ converge.

We say that the reverse Collatz sequence of $A$ is *trivial* if $A$ is the only odd number in its reverse Collatz sequence. We now record some properties of a reverse Collatz sequence given in Proposition 1 of [1].

**Proposition 3.1.** *Let $A$ be an odd number.*

1. *If $A$ is divisible by 3 then the reverse Collatz sequence of $A$ is trivial.*
2. *Let $p_i$ denote the subsequence of odd numbers in the reverse Collatz sequence of $A$. If $p_i \equiv 0 \ mod\ 3$, then $p_{i+1}$ do not exist. Otherwise, $p_{i+1}$ is the smallest odd number before $p_i$ in any Collatz sequence and*

$$p_{i+1} = \begin{cases} \frac{2p_i-1}{3} & \text{if } p_i \equiv 2 \ mod\ 3 \\ \\ \frac{4p_i-1}{3} & \text{if } p_i \equiv 1 \ mod\ 3 \end{cases} \tag{3}$$

3. *The only odd number that can be a jump from another odd number in a reverse Collatz sequence is the starting number of the reverse Collatz sequence.*

**Corollary 3.1.** *Let $A$ be an odd number and let $B = 2^k A + 1$ such that $k \geq 1$. Let $p_i$ denote the sequence of odd numbers in the reverse Collatz sequence of $p_0 = B$. Then*

$$p_1 = \begin{cases} 2^{k+2}(\frac{A}{3}) + 1 & \text{if } A \equiv 0 \ mod\ 3; \\ \\ 2(\frac{2^k A-1}{3}) + 1 & \text{if } A \not\equiv 0 \ mod\ 3. \end{cases}$$

*Proof.*

If $A \not\equiv 0 \bmod 3$, consider the four following situations: $k$ is even and $A \equiv 1 \bmod 3$ implies $B \equiv 2 \bmod 3$; $k$ is even and $A \equiv 2 \bmod 3$ implies $B \equiv 0 \bmod 3$; $k$ is odd and $A \equiv 1 \bmod 3$ implies $B \equiv 0 \bmod 3$; $k$ is odd and $A \equiv 2 \bmod 3$ implies $B \equiv 2 \bmod 3$.

By part (2) of Proposition 3.1, $p_1$ exists only if $B \not\equiv 0 \bmod 3$. Consequently, when $B \equiv 2 \bmod 3$, by part (2) of Proposition 3.1,

$$p_1 = \frac{2B-1}{3} = \frac{2^{k+1}A+1}{3} = \frac{2^{k+1}A-2+3}{3} = 2(\frac{2^k A-1}{3}) + 1.$$

Finally, if $A \equiv 0 \bmod 3$, then $B \equiv 1 \bmod 3$. Therefore, by part (2) of Proposition 3.1,

$$p_1 = \frac{4B-1}{3} = \frac{2^{k+2}A+3}{3} = 2^{k+2}(\frac{A}{3}) + 1.$$

□

**Theorem 3.1.** *Let $A$ be an odd number and let $p_i$, $q_i$, and $f_i$ denote the subsequence of odd numbers in the reverse Collatz sequence of $A$, $2A+1$, and $4A+1$, respectively.*

1. *If $A \equiv 0 \mod 3$, write $A = 3^n B$ such that $n \geq 1$, and $B$ is an odd number such that $B \not\equiv 0 \mod 3$. Then, for $i = 1, 2, \ldots, n$, $q_i = 2^{2i+1}(3^{n-i}B) + 1$ and $f_i = 2^{2i+2}(3^{n-i}B) + 1$. If $B \equiv 1 \mod 3$, then $q_n \equiv 0 \mod 3$. On the other hand, if $B \equiv 2 \mod 3$, then $f_n \equiv 0 \mod 3$.*

2. *Let $A \not\equiv 0 \mod 3$. Then the following conditions are true.*

   (a) *For some $k \geq 1$, $p_k \not\equiv 2 \mod 3$. If $p_k \equiv 1 \mod 3$, then $p_k = 3n+1$ and $p_{k+1} = 4n+1$ for some even number $n$.*

   (b) *If $A \equiv 1 \mod 3$, the reverse Collatz sequence of $2A+1$ is trivial. The reverse Collatz sequences of both $A$ and $4A+1$ are not trivial, and $f_i = 2p_i + 1$, for $1 \leq i \leq k$. If $p_k \equiv 0 \mod 3$, then $f_{k+1} = 2^3\left(\frac{p_k}{3}\right) + 1$. On the other hand, if $p_k \equiv 1 \mod 3$, then $f_k \equiv 0 \mod 3$.*

   (c) *Finally, if $A \equiv 2 \mod 3$, then the reverse Collatz sequence of $4A+1 \equiv 0$ is trivial. The reverse Collatz sequences of both $A$ and $2A+1$ are not trivial, and $q_i = 2p_i + 1$, for $0 \leq i \leq k$. If $p_k \equiv 0 \mod 3$, then $q_{k+1} = 2^3\left(\frac{p_k}{3}\right) + 1$. On the other hand, if $p_k \equiv 1 \mod 3$, then $q_k \equiv 0 \mod 3$.*

*Proof.*

1. If $A \equiv 0 \mod 3$, since $q_0 = 2A+1 \equiv 1 \mod 3$ and $f_0 = 4A+1 \equiv 1 \mod 3$, by Corollary 3.1, we get $q_1 = 2^3\left(\frac{A}{3}\right) + 1 = 2^3(3^{n-1}B) + 1$ and $f_1 = 2^4\left(\frac{A}{3}\right) + 1 = 2^4(3^{n-1}B) + 1$. Observe that, if $n > 1$, both $q_1$ and $f_1$ are equivalent to 1 mod 3. Hence, again, by Corollary 3.1, we get $q_2 = 2^5(3^{n-2}B) + 1$ and $f_1 = 2^6(3^{n-2}B) + 1$. Continuing the same argument, we get for $i = 1, 2, \ldots, n$, $q_i = 2^{2i+1}(3^{n-i}B) + 1$ and $f_i = 2^{2i+2}(3^{n-i}B) + 1$. Now, if $n$ is odd and $B \equiv 1 \mod 3$, then $q_n = 2^{2n+1}B + 1 \equiv 0 \mod 3$. If $n$ is even and if $B \equiv 2 \mod 3$, then $f_n = 2^{2n+2}B + 1 \equiv 0 \mod 3$.

2. Let $A \not\equiv 0 \mod 3$.

   (a) If $p_i \equiv 2 \mod 3$, then $p_{i+1} = (2p_i - 1)/3$, by part 2 of Proposition 3.1. Therefore $p_{i+1} < p_i$. If all the odd terms of the reverse Collatz sequence of $A$ are equivalent to 2 mod 3 then the sequence is strictly decreasing. The smallest such odd number that such a sequence can reach is 5, in which case, the next number in the sequence is 3, a contradiction. Hence, there is a $k$ such that $p_k \not\equiv 2 \mod 3$. If $p_k \equiv 1 \mod 3$, and since $p_k$ is odd, $p_k = 3n+1$ for some even number $n$. Moreover, by part 2 of Proposition 3.1, $p_{k+1} = (4p_k - 1)/3 = 4n+1$.

(b) By Part (1) of Proposition 3.1, the reverse Collatz sequence is trivial if the starting number is a multiple of 3. If $A \equiv 1 \bmod 3$, then $2A+1 \equiv 0 \bmod 3$ and $4A+1 \equiv 2 \bmod 3$. Therefore, the reverse Collatz sequence of $2A+1$ is trivial, whereas, the reverse Collatz sequences of both $A$ and $4A+1$ are not trivial. By part (2) of Proposition 3.1, $p_1 = (4A-1)/3$, and $f_1 = (2f_0-1)/3 = (2(4A+1)-1)/3 = (8A+1)/3$. Hence

$$2p_1 + 1 = 2\left(\frac{4A-1}{3}\right) + 1 = \frac{8A+1}{3} = f_1.$$

We use induction to prove that $f_i = 2p_i + 1$, if $p_i \equiv 2$, for $i > 1$. Assume the hypothesis is true for $i = r$, that is $p_r \equiv 2 \bmod 3$ and $f_r = 2p_r + 1$. Consequently, $f_r \equiv 2 \bmod 3$. Therefore, by part 2 of Proposition 3.1, $p_{r+1} = (2p_r - 1)/3$ and $f_{r+1} = (2f_r - 1)/3$. This implies

$$2p_{r+1} + 1 = 2\left(\frac{2p_r - 1}{3}\right) + 1 = \frac{4p_r + 1}{3} = \frac{2(2p_r + 1) - 1}{3} = \frac{2f_r - 1}{3} = f_{r+1}.$$

Therefore, it follows by induction that $f_i = 2p_i + 1$ for $1 \leq i \leq k$.

If $p_k \equiv 0 \bmod 3$, then since $f_k = 2p_k + 1$, we get by Corollary 3.1 that $f_{k+1} = 2^3\left(\frac{p_k}{3}\right) + 1$. Clearly, if $p_k \equiv 1 \bmod 3$, then $f_k \equiv 0 \bmod 3$.

(c) If $A \equiv 2 \bmod 3$, then $q_0 = 2A+1 \equiv 2 \bmod 3$ and $4A+1 \equiv 0 \bmod 3$. Therefore, the reverse Collatz sequence of $4A+1$ is trivial, whereas, the reverse Collatz sequences of both $A$ and $2A+1$ are not trivial. By part (2) of Proposition 3.1, $p_1 = (2A-1)/3$, and $q_1 = (2q_0 - 1)/3 = (2(2A+1) - 1)/3 = (4A+1)/3$. Consequently,

$$2p_1 + 1 = 2\left(\frac{2A-1}{3}\right) + 1 = \frac{4A+1}{3} = q_1.$$

We repeat the induction argument in part (2b) to get $q_i = 2p_i + 1$ for $0 \leq i \leq k$. Again, by Corollary 3.1, if $p_k \equiv 0 \bmod 3$, since $q_k = 2p_k + 1$, we get that $q_{k+1} = 2^3\left(\frac{p_k}{3}\right) + 1$. On the other hand, if $p_k \equiv 1 \bmod 3$, then $q_k \equiv 0 \bmod 3$.

□

# 4 Reduction of the Collatz conjecture to odd numbers that are congruent to 21 mod 24.

Let $A$ be an odd number. In this section, we show that the Collatz sequence of $A$ is equivalent to an integer of the form $24t + 21$ where $t \geq 0$.

**Theorem 4.1.** *Let $A > 1$ be an odd number and let $a_i$ denote the sequence of odd numbers in the Collatz sequence of $A$ with $a_0 = A$. Construct a sequence $b_i$ such that*

$$b_0 = \begin{cases} 12\left(2^4\left(\frac{A}{3}\right)+1\right)+9 & \text{if } A \equiv 0 \bmod 3; \\ 12\left(\frac{4A-1}{3}\right)+9 & \text{if } A \equiv 1 \bmod 3; \\ 12\left(\frac{2A-1}{3}\right)+9 & \text{if } A \equiv 2 \bmod 3. \end{cases}$$

*If for any $i$, $a_i = 1$, then $b_i = 12 \times 1 + 9 = 21$. Let $i \geq 1$ and let $a_{i-1} > 1$. If $a_{i-1} \equiv 3 \bmod 4$ then let $b_i = 12\left(\frac{a_{i-1}-1}{2}\right)+9$. If $a_{i-1} \equiv 1 \bmod 4$, then $a_{i-1} = 2^k N + 1$ for some odd number $N$, and $k \geq 2$. If $k > 2$, then $b_i = 12a_{i-1} + 9$. If $k = 2$, then $a_{i-1}$ is a jump from an odd number $P$, such that, $P$ is not a jump from an odd number. If $P = 1$, then $b_i = 12 \times 1 + 9 = 21$. If $P > 1$, write $P = 2^r q + 1$ such that $q$ is an odd number. Then, $r \neq 2$. If $r = 1$, let $b_i = 12\left(\frac{P-1}{2}\right)+9$. Otherwise, let $b_i = 12P + 9$.*

*Then, for all $i \geq 0$,*

1. *$b_i$ is odd, $b_i$ is divisible by 3, and $b_i \equiv 9 \bmod 12$.*
2. *the Collatz sequences of $a_i$ and $b_i$ are equivalent.*
3. *if $a_i > 1$, write $a_i = 2^k n_i + 1$ such that $k \geq 1$ and $n_i$ is an odd number. If $k \neq 2$, then*

$$b_{i+1} = \begin{cases} 3(2a_i+1), & \text{if } k = 1, \\ 3(4a_i+3), & \text{otherwise.} \end{cases}$$

*If $k = 2$, then $a_i$ is a jump from an odd number $j_i$, such that, $j_i$ is not a jump from an odd number. If $j_i > 1$, write $j_i = 2^r q_i + 1$ such that, $r \geq 1$ and $q_i$ is an odd number. Then $r \neq 2$, and*

$$b_{i+1} = \begin{cases} 3(2j_i+1), & \text{if } r = 1, \\ 3(4j_i+3), & \text{otherwise.} \end{cases}$$

*Proof.* 1. It follows by definition that $b_i$ is odd, $b_i$ is divisible by 3, and $b_i \equiv 9 \bmod 12$.

2. When $A \equiv 0 \bmod 3$, let $p_1$ denote the odd number that occurs after $4A + 1$ in the reverse Collatz sequence of $4A + 1$. Then by Corollary 3.1, $p_1 = 2^4\left(\frac{A}{3}\right)+1$. We see that

$$b_0 = 12p_1 + 9 = 12\left(2^4\left(\frac{A}{3}\right)+1\right)+9 = 4^3A + 4^2 + 4 + 1.$$

Thus, $b_0$ is a jump of height 3 from $A$.

Now, let $p_1$ denote the odd number that comes after $A$ in the reverse Collatz sequence of $A$. Then by Equation (3), we get

$$p_1 = \begin{cases} \frac{2A-1}{3}, & \text{if } A \equiv 2 \bmod 3 \\ \\ \frac{4A-1}{3}, & \text{if } A \equiv 1 \bmod 3 \end{cases}$$

Like before, we see that $12p_1 + 9 = b_0$ which is a jump of height of 2 from $A$, when $A \equiv 1 \bmod 3$, and a jump of height of 1 from $A$, when $A \equiv 2 \bmod 3$.

Now let $i \geq 1$. When $a_{i-1} \equiv 3 \bmod 4$, then $a_i = \frac{3a_{i-1}+1}{2}$. Consequently,

$$b_i = 12\left(\frac{a_{i-1}-1}{2}\right) + 9 = 6a_{i-1} + 3 = 4a_i + 1.$$

Thus, $b_i$ is a jump of height 1 from $a_i$.

Finally, let $a_{i-1} \equiv 1 \bmod 4$. If $k > 2$, $a_i = \frac{3a_{i-1}+1}{4}$.

Then $4^2 a_i + 4 + 1 = 12a_{i-1} + 9 = b_i$ is a jump from $a_i$ of height 2.

When $k = 2$ the Collatz sequence of $a_{i-1}$ is equivalent to the Collatz sequence of $P$. By the definition of $P$, $r \neq 2$. If $r > 2$, then $a_i = (3P+1)/4$, and if $r = 1$, $a_i = (3P+1)/2$. We repeat the above argument and get that $b_i$ is a jump of height 2 from $a_i$ when when $k > 2$, and a jump of height 1, otherwise.

Thus, for any $i$, $b_i$ is a jump from $a_i$. Hence, the Collatz sequences of $a_i$ and $b_i$ are equivalent.

3. Since $a_i$ is odd, when $k = 1$, $a_i \equiv 3 \bmod 4$. Therefore

$$b_{i+1} = 12\left(\frac{a_i - 1}{2}\right) + 9 = 3(2a_i + 1).$$

When $k > 2$, $a_i \equiv 1 \bmod 4$. Therefore, $b_{i+1} = 12a_i + 9 = 3(4a_i + 3)$.

Let $k = 2$, then $r \neq 2$ by definition of $j_i$, and

$$b_{i+1} = \begin{cases} 12\left(\frac{j_i-1}{2}\right) + 9 = 3(2j_i + 1), & \text{if } r = 1, \\ \\ 12j_i + 9 = 3(4j_i + 3), & \text{otherwise.} \end{cases}$$

□

**Corollary 4.1.** *1. Given an odd number $a$, the Collatz sequence of $a$ is equivalent to an integer of the form $24t + 21$ where $t \geq 0$.*

*2. Consider a number of the form $24t + 21$, where $t \geq 0$. Let $p = 2t + 1$. If $t = 0$, the the Collatz sequences of $24t + 21$ and 1 merge. If $t > 0$, write $p = 2^k n + 1$, where $k \geq 1$, and $n$ is an odd number. If $k > 2$, the Collatz sequences of $24t + 21$ and $p$ merge. Otherwise, the Collatz sequences of $24t + 21$ and $2p + 1$ merge.*

*Proof.* 1. Applying Theorem 4.1, we get that the Collatz sequence of $a$ is equivalent to the Collatz sequence of an integer of the form $12d + 9$ where $d$ is an odd number. Write $d = 2t + 1$ such that $t \geq 0$. This gives us the desired result.

2. This result is a direct application of Theorem 4.1.

□

**Example 4.1.** Let $a_i$ denote the odd numbers in the Collatz sequence of 319, with $a_0 = 319$. For each $i$, if $a_i > 1$, write $a_i = 2^k n_i + 1$, where $n_i$ is an odd number. If $k > 2$ then $e_i = 4a_i + 3$. If $k = 1$, then $d_i = 2a_i + 1$. If $k = 2$, then let $p_i$ be an odd number such that, $p_i$ is not a jump from an odd number. If $p_i = 1$, then $e_i = 4 \times 1 + 3 = 7$. If $p_i > 1$, write $p_i = 2^r q_i + 1$ such that $q_i$ is an odd number. Then, if $r = 1$, then $d_i = 2q_i + 1$ and if $r > 1$ then $e_i = 4p_i + 3$. Let $b_i$ be defined as in Theorem 4.1. We get the following table from the Collatz sequence of 319.

| $b_i$ | $a_i$ | $d_i$ | $3d_i$ | $e_i$ | $3e_i$ |
|---|---|---|---|---|---|
| $5109 = 12 \times 425 + 9$ | $319 = 2 \times 159 + 1$ | 639 | 1917 | | |
| $1917 = 12 \times 159 + 9$ | $479 = 2 \times 239 + 1$ | 959 | 2877 | | |
| $2877 = 12 \times 239 + 9$ | $719 = 2 \times 359 + 1$ | 1439 | 4317 | | |
| $4317 = 12 \times 359 + 9$ | $1079 = 2 \times 539 + 1$ | 2159 | 6477 | | |
| $6477 = 12 \times 539 + 9$ | $1619 = 2 \times 809 + 1$ | 3239 | 9717 | | |
| $9717 = 12 \times 809 + 9$ | $2429 = 4 \times 607 + 1$ | 1215 | 3645 | | |
| $3645 = 12 \times 303 + 9$ | $911 = 2 \times 455 + 1$ | 1823 | 5469 | | |
| $5469 = 12 \times 455 + 9$ | $1367 = 2 \times 683 + 1$ | 2735 | 8205 | | |
| $8205 = 12 \times 683 + 9$ | $2051 = 2 \times 1025 + 1$ | 4103 | 12309 | | |
| $12309 = 12 \times 1025 + 9$ | $3077 = 4 \times 769 + 1$ | | | 3079 | 9237 |
| $9237 = 12 \times 769 + 9$ | $577 = 2^6 \times 9 + 1$ | | | 2311 | 6933 |
| $6933 = 12 \times 577 + 9$ | $433 = 2^4 \times 27 + 1$ | | | 1735 | 5205 |
| $5205 = 12 \times 433 + 1$ | $325 = 4 \times 81 + 1$ | | | 327 | 981 |
| $981 = 12 \times 81 + 9$ | $61 = 4 \times 15 + 1$ | 31 | 93 | | |
| $93 = 12 \times 7 + 9$ | $23 = 2 \times 11 + 1$ | 47 | 141 | | |
| $141 = 12 \times 11 + 9$ | $35 = 2 \times 17 + 1$ | 71 | 213 | | |
| $213 = 12 \times 17 + 9$ | $53 = 4^2 \times 3 + 4 + 1$ | 7 | 21 | | |
| $21 = 12 \times 1 + 9$ | $5 = 4 \times 1 + 1$ | | | 7 | 21 |
| $21 = 12 \times 1 + 9$ | 1 | | | | |

# 5 Network of Collatz sequences.

In this section, we construct a network of diagonal arrays which contain the Collatz sequences of all odd numbers.

**Theorem 5.1.** *For $n \not\equiv 1 \bmod 3$, define a diagonal array as follows. Let $u_0 = 4n + 1$ and $u_i = 2u_{i-1} + 1$, for $i \geq 1$. For $j \geq 0$, let $v_{0,j} = u_j$, and for $k \geq 1$, let $v_{k,k} = 3v_{k-1,k-1} + 2$. Finally, for $j > i$, let $v_{i,j} = 2v_{i,j-1} + 1$. We get an array*

$$\begin{array}{cccccccccc} u_0 & u_1 & u_2 & u_3 & u_4 & u_5 & u_6 & u_7 & \dots \\ & v_{1,1} & v_{1,2} & v_{1,3} & v_{1,4} & v_{1,5} & v_{1,6} & v_{1,7} & \dots \\ & & v_{2,2} & v_{2,3} & v_{2,4} & v_{2,5} & v_{2,6} & v_{2,7} & \dots \\ & & & v_{3,3} & v_{3,4} & v_{3,5} & v_{3,6} & v_{3,7} & \dots \\ & & & & \dots & \dots & & & \end{array}$$

*with the following properties:*

1. $u_k \not\equiv 2 \mod 3$ *for all* $k \geq 0$, *whereas,* $v_{i,j} \equiv 2 \mod 3$, *if* $i > 0$ *and* $j > 0$.
2. $u_0 \equiv 1 \mod 4$ *and* $v_{i,i} \equiv 1 \mod 4$ *for all* $i$. $u_i \not\equiv 1 \mod 4$ *for* $i > 0$ *and* $v_{i,j} \not\equiv 1 \mod 4$ *if* $i \neq j$.
3. *For* $i \geq 0$, *if* $u_i \equiv 1 \mod 3$ *then* $u_{i+1} \equiv 0 \mod 3$. *On the other hand, if* $u_i \equiv 0 \mod 3$ *then* $u_{i+1} \equiv 1 \mod 3$.
4. *If* $u_0 \equiv 0 \mod 3$, *then write* $u_0 = 3N$ *for some* $N$. *For* $i \geq 0$, *let* $j_i$ *denote the sequence of jumps of height* $i$ *from* $N$, *then* $u_{2i} = 3j_i$ *and* $u_{2i+1} = 6j_i + 1$. *Alternately, if* $u_1 \equiv 0 \mod 3$, *then write* $u_1 = 3N$ *for some* $N$. *For* $i \geq 0$, *let* $j_i$ *denote the sequence of jumps of height* $i$ *from* $N$. *Then,* $u_{2i+1} = 3j_i$ *and* $u_{2i} = 6j_i + 1$.
5. *For* $j \geq 1$, *the* $j$*-th column is the first few odd numbers in the Collatz sequence of* $u_j$.
6. *For* $j \geq i > 0$, $v_{i,j} = 3v_{i-1,j-1} + 2$.
7. *Let* $a_{i,i}$ *denote the odd number that comes after* $v_{i,i}$ *in the Collatz sequence of* $v_{i,i}$. *If* $n$ *is even, then for* $k \geq 0$, $a_{2k,2k} \equiv 1 \mod 3$ *and* $v_{2k+1,2k+1} = 4a_{2k,2k} + 1$. *Moreover,* $a_{2k,2k}$ *is smaller than both* $v_{2k,2k}$ *and* $v_{2k+1,2k+1}$. *If* $n$ *is odd, then for* $k \geq 1$, $a_{2k-1,2k-1} \equiv 1 \mod 3$ *and* $v_{2k,2k} = 4a_{2k-1,2k-1} + 1$. *Like before,* $a_{2k-1,2k-1}$ *is smaller than both* $v_{2k-1,2k-1}$ *and* $v_{2k,2k}$.
8. *If* $n$ *is odd, then for* $k \geq 1$, *the Collatz sequences of* $u_{2k-1}$ *and* $u_{2k}$ *merge. If* $n$ *is even, then for* $k \geq 0$, *the Collatz sequences of* $u_{2k}$ *and* $u_{2k+1}$ *merge.*
9. *For every* $n > 0$, *such that,* $n \not\equiv 1 \mod 3$, *the Collatz sequence of* $u_0$ *merges with the Collatz sequence of a smaller number* $a$ *such that* $a \not\equiv 2 \mod 3$.

*Proof.* 1. Since $n \not\equiv 1 \mod 3$, $u_0 = 4n + 1 \not\equiv 2 \mod 3$. If $u_i \not\equiv 2 \mod 3$, then $u_{i+1} = 2u_i + 1 \not\equiv 2 \mod 3$. Consequently, $u_k \not\equiv 2 \mod 3$ for all $k \geq 0$.

For $i > 0$, $v_{i,i} \equiv 2 \mod 3$, by definition. If $v_{i,j} \equiv 2 \mod 3$, then $v_{i,j+1} = 2v_{i,j} + 1 \equiv 2 \mod 3$. Thus, it follows that $v_{i,j} \equiv 2 \mod 3$, if $i > 0$ and $j > 0$.

2. $u_0 \equiv 1 \mod 4$ by definition. $v_{i,i} = 3v_{i-1,i-1} + 2 \equiv 1 \mod 4$, if $v_{i-1,i-1} \equiv 1 \mod 4$. Since $v_{0,0} = u_0$, it follows that $v_{i,i} \equiv 1 \mod 4$ for $i \geq 0$. For $i > 0$, $u_i = 2u_{i-1} + 1 \not\equiv 1 \mod 4$ since $u_{i-1}$ is an odd number. Similarly, $v_{i,j} = 2v_{i,j-1} + 1 \not\equiv 1 \mod 4$, when $i \neq j$.

3. Since $u_{i+1} = 2u_i + 1$, if $u_i \equiv 0 \mod 3$, then $u_{i+1} \equiv 1 \mod 3$. Similarly, if $u_i \equiv 1 \mod 3$, then $u_{i+1} \equiv 0 \mod 3$

4. We prove this result by using the induction principle. If $u_0 = 3N$, then $u_1 = 6N + 1$. Thus $u_0 = 3j_0$ and $u_1 = 6j_0 + 1$. Assume $u_{2i} = 3j_i$ and $u_{2i+1} = 6j_i + 1$. Then, $u_{2(i+1)=2i+2} = 2u_{2i+1} + 1 = 3(4j_i + 1) = 3j_{i+1}$. And $u_{2(i+1)+1=2i+3} = 2u_{2i+2} + 1 = 6j_{i+1} + 1$. Hence the result. The proof is similar for the case when $u_1 \equiv 0 \mod 3$.

5. When $j \geq 1$, $v_{0,j} = u_j$. We know from Part 2 that $v_{i,j} \equiv 3 \bmod 4$ if $i \neq j$. Therefore, the odd number that comes after $v_{i,j}$ in the Collatz sequence of $v_{i,j}$ is $(3v_{i,j}+1)/2$. We do not compute the odd numbers that come after $v_{i,i}$ in this part. Hence, assume that $j > i$.

   For $k \geq 1$, let $z_k$ be defined as follows:

$$z_k = 2^k + 2^{k-1} + 2^{k-2} + \cdots + 2 + 1 = \sum_{i=0}^{k} 2^i = \frac{2^{k+1}-1}{2-1} = 2^{k+1} - 1.$$

   Then, by definition, for $j > i$, $v_{i,j} = 2^{j-i}v_{ii} + z_{j-i-1}$, and

$$v_{i+1,j} = 2^{j-i-1}v_{i+1,i+1} + z_{j-i-2} = 2^{j-i-1}(3v_{i,i}+2) + z_{j-i-2}.$$

$$\frac{3v_{i,j}+1}{2} = \frac{3 \times 2^{j-i}v_{ii} + 3 \times z_{j-i-1} + 1}{2} = 3 \times 2^{j-i-1}v_{ii} + \frac{3z_{j-i-1}+1}{2}.$$

$$\frac{3z_{j-i-1}+1}{2} = \frac{3(2^{j-i}-1)+1}{2} = 3\times 2^{j-i-1} - 1 = 2\times 2^{j-i-1} + (2^{j-i-1}-1) = 2^{j-i} + z_{j-i-2}.$$

   Consequently,

$$\frac{3v_{i,j}+1}{2} = v_{i+1,j}.$$

   This implies $v_{i+1,j}$ is the odd number that comes after $v_{i,j}$ in the Collatz sequence of $v_{i,j}$. Hence, for $j \geq 1$, the $j$-th column is the first few odd numbers in the Collatz sequence of $u_j$.

6. For $j \geq i > 0$, when $i = j$, we have that $v_{i,j} = 3v_{i-1,j-1} + 2$, by definition. Let $i \neq j$, then $v_{i,j} = 2v_{i,j-1} + 1$. But $v_{i-1,j-1} \not\equiv 1 \bmod 4$, when $i \neq j$. Hence $v_{i,j-1} = (3v_{i-1,j-1}+1)/2$. Consequently, $v_{i,j} = 3v_{i-1,j-1} + 2$.

7. If $n$ is even, then $v_{0,0} = 2^k n + 1$ such $k > 2$. By Part 3b of Proposition 2.1, $a_{0,0} = 2^{k-2}3n + 1 \equiv 1 \bmod 3$. Moreover, $a_{0,0} = (3v_{0,0}+1)/4$. Therefore, by Theorem 2.1, $v_{1,1} = 4a_{0,0} + 1$. $a_{0,0}$ is smaller than both $v_{0,0}$ and $v_{1,1}$. Now the proof proceeds by induction. Assume for $k \geq 0$, $a_{2k,2k} \equiv 1 \bmod 3$, $v_{2k+1,2k+1} = 4a_{2k,2k} + 1$, and $a_{2k,2k}$ is smaller than both $v_{2k,2k}$ and $v_{2k+1,2k+1}$. Since $a_{2k,2k}$ is odd, by part 3 of Proposition 2.1, $a_{2k+1,2k+1} = (3v_{2k+1,2k+1}+1)/2^j$ where $j \geq 3$. By Theorem 2.1, $v_{2k+2,2k+2} = 2^j a_{2k+1,2k+1} + 1$. Hence, $a_{2k+2,2k+2} = (3v_{2k+2,2k+2}+1)/4$. Again, by Theorem 2.1, $v_{2k+3,2k+3} = 4a_{2k+2,2k+2} + 1$. We see that $a_{2k+2,2k+2}$ is smaller that $v_{2k+2,2k+2}$ and $v_{2k+1,2k+1}$. Hence, the proof.

   When $n$ is odd, by part 3 of Proposition 2.1, $a_{0,0} = (3v_{0,0}+1)/2^r$, such that $r \geq 3$. Therefore, by Theorem 2.1, $v_{1,1} = 2^r a_{0,0} + 1$. Hence $a_{1,1} = (3v_{1,1}+1)/4$ which implies $v_{2,2} = 4a_{1,1} + 1$. Moreover $a_{1,1} = 2^{r-2}3a_{0,0} + 1 \equiv 1 \bmod 3$. We also see that $a_{1,1}$ is smaller than $v_{1,1}$ and $v_{2,2}$. Now, we apply a similar argument, as in the $n$ is even case, to show that for $k \geq 1$, $a_{2k-1,2k-1} \equiv 1 \bmod 3$, $v_{2k,2k} = 4a_{2k-1,2k-1} + 1$, and $a_{2k-1,2k-1}$ is smaller than both $v_{2k-1,2k-1}$ and $v_{2k,2k}$

8. By Part 7 and Part 1 of Proposition 2.1, we get that If $n$ is odd, then for $k \geq 1$, the Collatz sequences of $v_{2k-1,2k-1}$ and $v_{2k,2k}$ are equivalent; and if $n$ is even then for $k \geq 0$, the Collatz sequences of $v_{2k,2k}$ and $v_{2k+1,2k+1}$ are equivalent. Consequently, it follows that if $n$ is odd for $k \geq 1$, the Collatz sequences of $u_{2k-1}$ and $u_{2k}$ merge; and if $n$ is even, then for $k \geq 0$, the Collatz sequences of $u_{2k}$ and $u_{2k+1}$ merge.

9. If $n$ is odd, then by Part 1 of Proposition 2.1, we get that the Collatz sequences of $u_0 = 4n + 1$ and $n$ are equivalent. If $n \equiv 0 \bmod 3$, since $n < u_0$, in this case, the Collatz sequence of $u_0$ merges with the Collatz sequence of a smaller number. Now, let $n \equiv 2 \bmod 3$. By Part 2a of Theorem 3.1, the reverse Collatz sequence of $n$ reaches a number $a$ such that $a \not\equiv 2 \bmod 3$. By definition of the reverse Collatz sequence, it is strictly decreasing till it reaches $a$. Hence $a < u_0$. Finally, if $n$ is even, we get that the Collatz sequence of $u_0$ reaches a smaller number congruent to 1 mod 3 by Part 7.

□

**Example 5.1.** Let $u_i$ and $v_{i,j}$ be defined as in Theorem 5.1. Let $t_i$ denote the odd number congruent to 1 mod 3 such that the Collatz sequences of $v_{i,i}$ and $t_i$ merge:

$n = 0$ :

| | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| $u_i$ : | 1 | 3 | 7 | 15 | 31 | 63 | 127 | 255 | 511 | 1023 | 2047 | 4095 | 8191 |
| $v_{1,i}$ : | | 5 | 11 | 23 | 47 | 95 | 191 | 383 | 767 | 1535 | 3071 | 6143 | 12287 |
| $v_{2,i}$ : | | | 17 | 35 | 71 | 143 | 287 | 575 | 1151 | 2303 | 4607 | 9215 | 18431 |
| $v_{3,i}$ : | | | | 53 | 107 | 215 | 431 | 863 | 1727 | 3455 | 6911 | 13823 | 27647 |
| $v_{4,i}$ : | | | | | 161 | 323 | 647 | 1295 | 2591 | 5183 | 10367 | 20735 | 41471 |
| $v_{5,i}$ | | | | | | 485 | 971 | 1943 | 3887 | 7775 | 15551 | 31103 | 62207 |
| $v_{6,i}$ : | | | | | | | 1457 | 2915 | 5831 | 11663 | 23327 | 46655 | 93311 |
| $v_{7,i}$ : | | | | | | | | 4373 | 8747 | 17495 | 34991 | 69983 | 139967 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| $v_{i,i}$ : | 1 | 5 | 17 | 53 | 161 | 485 | 1457 | 4373 | ... |
| $t_i$ : | | 1 | 1 | 13 | 13 | 121 | 121 | 1093 | 1093 |

$n = 3$ :

| | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| $u_i$ : | 13 | 27 | 55 | 111 | 223 | 447 | 895 | 1791 | 3583 | 7167 | 14335 | 28671 |
| $v_{1,i}$ : | | 41 | 83 | 167 | 335 | 671 | 1343 | 2687 | 5375 | 10751 | 21503 | 43007 |
| $v_{2,i}$ : | | | 125 | 251 | 503 | 1007 | 2015 | 4031 | 8063 | 16127 | 32255 | 64511 |
| $v_{3,i}$ : | | | | 377 | 755 | 1511 | 3023 | 6047 | 12095 | 24191 | 48383 | 96767 |
| $v_{4,i}$ : | | | | | 1133 | 2267 | 4535 | 9071 | 18143 | 36287 | 72575 | 145151 |
| $v_{5,i}$ : | | | | | | 3401 | 6803 | 13607 | 27215 | 54431 | 108863 | 217727 |
| $v_{6,i}$ : | | | | | | | 10205 | 20411 | 40823 | 81647 | 163295 | 326591 |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| $v_{i,i}$ : | 13 | 41 | 125 | 377 | 1133 | 3401 | 10205 | ... |
| $t_i$ : | 13 | 31 | 31 | 283 | 283 | 2551 | 2551 | |

**Corollary 5.1.** *Every odd number occurs in one of the diagonal arrays defined in Theorem 5.1.*

*Proof.* Given an odd number $u_i \not\equiv 2 \bmod 3$, let $u_{i-1} = (u_i - 1)/2$. This sequence eventually reaches a number that is congruent to 1 mod 4, because it is strictly decreasing. Moreover, $u_i \not\equiv 2 \bmod 3$ for all $i$. Thus, every $u_i \not\equiv 2 \bmod 3$ occur in one of the diagonal arrays defined in Theorem 5.1. Now consider an odd integer $a \equiv 2 \bmod 3$. By Part 2a of Theorem 3.1, the reverse Collatz sequence of $a$ reaches a number that is not congruent to 2 mod 3. Hence, the result. □

# References


[1] Ahmed Maya M, *A window to the convergence of the Collatz sequence* (to appear in Integers Journal), arXiv:1510.01274 [math.GM].

[2] Jeffrey C. Lagarias, *The 3x+1 Problem: An Annotated Bibliography, II (2000-2009)*, arXiv:math/0608208v5 [math.NT]

[3] Jeffrey C. Lagarias, *The 3x+1 problem and its generalizations*, Amer. Math. Monthly 92 (1985) 3-23.

[4] Wirsching, Günther J., *The Dynamical System Generated by the 3n+1 Function*, Lecture Notes in Mathematics, Vol. 1681, Springer Verlag, 1998.